

134400-1



IN THE SPECIFICATION

Please amend the title as follows:

COLORED POLYMERIC RESIN COMPOSITION CONTAINING 1,8-DIAMINOANTHRAQUINONE DERIVATIVE, ARTICLE MADE THEREFROM, AND METHOD FOR MAKING THE SAME

Please amend the Abstract as follows:

In one embodiment, a colored polymeric resin composition, comprises: a polymeric resin; and a 1,8-diaminoanthraquinone derivative having a purity of greater than or equal to about 90 wt% and having a Formula (VIII):




134400-1


wherein  $R_2 - R_7$  are, individually, selected from the group consisting of a hydrogen atom, a hydroxyl group, an alkoxy group, an aryloxy group, an aliphatic group, an aromatic group, a heterocyclic group, a halogen atom, a cyano group, a nitro group,  $-\text{COR}_9$ ,  $-\text{COOR}_9$ ,  $-\text{NR}_9\text{R}_{10}$ ,  $-\text{NR}_{10}\text{COR}_{11}$ ,  $-\text{NR}_{10}\text{SO}_2\text{R}_{11}$ ,  $-\text{CONR}_9\text{R}_{10}$ ,  $-\text{CONHSO}_2\text{R}_{11}$ , and  $-\text{SO}_2\text{NHCOR}_{11}$ ; in which  $R_9$  and  $R_{10}$  are, individually, selected from the group consisting of a hydrogen atom, an aliphatic group, an aromatic group, and a heterocyclic group; wherein  $R_{11}$  is selected from the group consisting of an aliphatic group, an aromatic group, and a heterocyclic group; and wherein  $R$  is selected from the group consisting of hydrogen, an alkyl group containing 1 to 20 carbon atoms, a cycloalkyl group containing 3 to 20 carbon atoms, an allyl group containing 3 to 20 carbon atoms, a hydroxyl group, a 5-membered heterocyclic ring, and a 6-membered heterocyclic ring.

Please amend paragraph [0004] on page 2 of the specification as follows:

[0004] Disclosed herein are embodiments of colored polymeric resins, articles made therefrom, and methods for making the same. In one embodiment, a colored polymeric resin composition comprises: a polymeric resin and a 1,8-anthraquinone derivative having a purity of greater than or equal to about 90 wt% and having a Formula (~~XI~~ VIII):



134400-1




wherein  $R_2 - R_7$  are, individually, selected from the group consisting of a hydrogen atom, a hydroxyl group, an alkoxy group, an aryloxy group, an aliphatic group, an aromatic group, a heterocyclic group, a halogen atom, a cyano group, a nitro group,  $-\text{COR}_9$ ,  $-\text{COOR}_9$ ,  $-\text{NR}_9\text{R}_{10}$ ,  $-\text{NR}_{10}\text{COR}_{11}$ ,  $-\text{NR}_{10}\text{SO}_2\text{R}_{11}$ ,  $-\text{CONR}_9\text{R}_{10}$ ,  $-\text{CONHSO}_2\text{R}_{11}$ , and  $-\text{SO}_2\text{NHCOR}_{11}$ ; in which  $R_9$  and  $R_{10}$  are, individually, selected from the group consisting of a hydrogen atom, an aliphatic group, an aromatic group, and a heterocyclic group; wherein  $R_{11}$  is selected from the group consisting of an aliphatic group, an aromatic group, and a heterocyclic group; and wherein  $R$  is selected from the group consisting of hydrogen, an alkyl group containing 1 to 20 carbon atoms, a cycloalkyl group containing 3 to 20 carbon atoms, an allyl group containing 3 to 20 carbon atoms, a hydroxyl group, a 5-membered heterocyclic ring, and a 6-membered heterocyclic ring.


Please amend paragraph [0005], which spans pages 2-3 of the specification, as follows:

[0005]In another embodiment, the colored polymeric resin composition comprises: a polymeric resin; and a 1,8-anthraquinone derivative having a Formula (~~IX~~ VIII):

134400-1



(XI)




(VIII)

wherein  $R_2 - R_7$  are, individually, selected from the group consisting of a hydrogen atom, a hydroxyl group, an alkoxy group, an aryloxy group, an aliphatic group, an aromatic group, a heterocyclic group, a halogen atom, a cyano group, a nitro group,  $-\text{COR}_9$ ,  $-\text{COOR}_9$ ,  $-\text{NR}_9\text{R}_{10}$ ,  $-\text{NR}_{10}\text{COR}_{11}$ ,  $-\text{NR}_{10}\text{SO}_2\text{R}_{11}$ ,  $-\text{CONR}_9\text{R}_{10}$ ,  $-\text{CONHSO}_2\text{R}_{11}$ , and  $-\text{SO}_2\text{NHCOR}_{11}$ ; in which  $R_9$  and  $R_{10}$  are, individually, selected from the group consisting of a hydrogen atom, an aliphatic group, an aromatic group, and a heterocyclic group; wherein  $R_{11}$  is selected from the group consisting of an aliphatic group, an aromatic group, and a heterocyclic group; and wherein  $R$  is selected from the group consisting of hydrogen, an alkyl group containing 1 to 20 carbon atoms, a cycloalkyl group containing 3 to 20 carbon atoms, an allyl group containing 3 to 20 carbon atoms, a hydroxyl group, a 5-membered heterocyclic ring, and a 6-membered heterocyclic ring. An article formed from this composition has a hue angle value of less than or equal to about 330 degrees (when used at a loading of 0.01 pph at an article thickness of 3.2 mm).


134400-1

Please amend paragraph [0036], which spans pages 14-15 of the specification, as follows:

[0036]The anthraquinone derivatives may be represented by the Formula (VIII):



(VIII)



(VIII)

wherein  $R_2 - R_7$  are, individually, selected from the group consisting of a hydrogen atom, a hydroxyl group, an alkoxy group, an aryloxy group, an aliphatic group, an aromatic group, a heterocyclic group, a halogen atom, a cyano group, a nitro group,  $-\text{COR}_9$ ,  $-\text{COOR}_9$ ,  $-\text{SR}_9$ ,  $-\text{NR}_9\text{R}_{10}$ ,  $-\text{NR}_{10}\text{COR}_{11}$ ,  $-\text{NR}_{10}\text{SO}_2\text{R}_{11}$ ,  $-\text{CONR}_9\text{R}_{10}$ ,  $-\text{CONHSO}_2\text{R}_{11}$ , and  $-\text{SO}_2\text{NHCOR}_{11}$ ; in which  $R_9$  and  $R_{10}$  are, individually, selected from the group consisting of a hydrogen atom, an aliphatic group, an aromatic group, and a heterocyclic group; wherein  $R_{11}$  is selected from the group consisting of an aliphatic group, an aromatic group, and a heterocyclic group, while  $R$  is hydrogen, an alkyl group containing 1 to 20 carbon atoms (e.g., methyl, ethyl, n-butyl, isopropyl, 2-ethylhexyl, n-decyl, n-

134400-1

octadecyl), a cycloalkyl group containing 3 to 20 carbon atoms (e.g., cyclopentyl, cyclohexyl), an allyl group containing 3 to 20 carbon atoms, which may contain a substituent(s) including, for example, a halogen atom (e.g., F, Cl, Br, I), a 5- or 6-membered heterocyclic ring, or the like. Some examples of the above 1,8-anthraquinone derivatives include: 1,8-bis(cyclohexylamino) anthraquinone; 1,8-bis(isopropylamino) anthraquinone; 1,8-bis(2-ethylhexylamino) anthraquinone; 1,8-bis(N,N-diethylamino-propylamino) anthraquinone; and 1,8-bis(N,N-diethylamino-ethylamino) anthraquinone, and their derivatives, and the like, as well as reaction products and combinations comprising at least one of the foregoing, with 1,8-bis(cyclohexylamino) anthraquinone preferred.

Please amend Table 4 of the specification as follows:

134400-1

TABLE 4					
Series	Chemical Structure	Ex. #	Structural details		
			R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>
1,5-dialkylamino anthraquinone		N/A	cyclohexyl	H	H
		1	<sup>2</sup> i-Pr	H	H
		2	N,N-diethylaminoethyl	H	H
1,8-dialkylamino anthraquinone		3	cyclohexyl	H	H
		4	i-Pr	H	H
		5	2-ethyl hexyl	H	H
		6	3-N,N-dimethylamino propylamine	H	H
		7	N,N-diethylaminoethyl	H	H

<sup>1</sup>Ex.# = example number<sup>2</sup>iso-prop